

Asymptotic behaviour of the weighted trace of the Schrödinger operator

Fatih Taşçi and Mehmet Bayramoğlu

Abstract

This paper is devoted to the study of the asymptotic behaviour of the weighted trace of the Schrödinger operator with operator coefficient.

M.S.C. 2000: 45M05, 39B42.

Key words: differential operators, weighted trace, asymptotics.

§1. Introduction

Let H be a separable Hilbert space with the inner product $\langle \cdot, \cdot \rangle$. In the Hilbert space $H_1 = L_2(\mathbb{R}_n^+; H)$ we consider the differential operator L generated by the expression

$$(1.1) \quad -\Delta u + Q(x)u, \quad x \in \mathbb{R}_n^+$$

and the boundary condition

$$(1.2) \quad u(x_1, x_2, \dots, x_n)|_{x_n=0} = 0,$$

where we denote by H_1 the Hilbert space of H valued square integrable functions with the scalar product

$$(1.3) \quad (f, g) = \int_{\mathbb{R}_n^+} (f(x), g(x))_H dx, \quad f(x), g(x) \in H_1.$$

Denote by \mathbb{R}_n^+ the set of points $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_n$, $x_n \geq 0$. Here $Q(x)$ is a self-adjoint operator acting on H . We shall assume that $Q(x)$ satisfies the following conditions:

(1⁰) The domain of definition $D(Q(x)) = D$ of $Q(x)$ is independent of x and $\overline{D} = H$ and $Q^*(x) = Q(x) \geq I$, where I is the identity operator on H . It is also assumed that $Q^{-1}(x)$ is completely continuous operator on H for each $x \in \mathbb{R}_n^+$.

(2⁰) $Q^{-1}(x) \in \sigma_1$, $\forall x \in \mathbb{R}_n^+$ and $\int_{\mathbb{R}_n^+} \|Q^{-k}(x)\|_1 dx < \infty$ for any $k > 0$, where σ_1 is the space of trace class operators and $\|\cdot\|_1$ is the norm in σ_1 [5].

$$(3^0) \left\| Q^\gamma(s) e^{-ctQ(s)} \right\|_1 \leq C \left\| Q^\gamma(x) e^{-f(c)tQ(x)} \right\|_1,$$

$$\left\| Q^\beta(x) Q^{-\beta}(s) \right\| \leq C, |x-s| \leq 1,$$

where $\gamma = 0, \beta = f(c) > 0$ and β is a real number.

$$(4^0) \left\| Q^\gamma(x) [Q(s) - Q(x)] Q^{-\gamma-b}(s) \right\| < C |x-s|,$$

$$|x-s| \leq 1, \quad 0 < b < 3/2.$$

(5⁰) $\int_{\mathbb{R}_n^+} Tr [Q^\gamma(x) e^{-qtQ(x)}] dx = O(1) \int_{\mathbb{R}_n^+} Tr [Q^\gamma(x) e^{-tQ(x)}] dx, \quad t \rightarrow 0$, where q is a positive constant.

(6⁰) Let $\alpha_1(x) \leq \alpha_2(x) \leq \dots \leq \alpha_n(x) \leq \dots$ be eigenvalues of the operator $Q(x)$. Denote

$$\rho_\beta(\lambda) = \frac{1}{2^n \pi^{\frac{n}{2}} \Gamma(\frac{n}{2} + 1)} \sum_j \int_{\alpha_j(x) \leq \lambda} \alpha_j^\beta(x) [\lambda - \alpha_j(x)]^{n/2} dx,$$

and assume that $\lim_{\lambda \rightarrow \infty} \rho_\beta(\lambda) = \infty$ and for large λ , the condition $\lambda \rho'_\beta(\lambda) < a \rho_\beta(\lambda)$ is satisfied for any positive number a . In this paper we used one and the same symbol C in order to denote any constant which may differ from each other.

Let $\varphi_k(x)$'s be functions that have compact support, continuous second order derivative and satisfying $\varphi_k(x)|_{x_n=0} = 0, f_k \in D$. Let us denote by D' the summation $\sum_{k=1}^m \varphi_k(x) f_k$. In the Hilbert space H_1 we consider the operator L' generated by $-\Delta u + Q(x)u$ with the domain of definition D' . This operator is symmetric and positive definite. The closure L of the operator L' is self adjoint.

The spectrum of the operator L generated by the expression (1.1) and by the boundary condition (1.2) consists of eigenvalues. Let $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \leq \dots$ be the eigenvalues of this operator, and $\psi_1(x), \psi_2(x), \dots, \psi_n(x), \dots$ the corresponding complete vector-valued orthonormal eigenfunctions.

The aim of this paper is to find asymptotic behaviour of $N_\beta(\lambda)$ as $\lambda \rightarrow \infty$. We define $N_\beta(\lambda)$ as follows

$$(1.4) \quad N_\beta(\lambda) = \sum_{\lambda_k < \lambda} \int_{\mathbb{R}_n^+} (Q^\beta(x) \psi_k(x), \psi_k(x)) dx,$$

and it is called weighted-trace of the operator L , where β is a real constant.

§2. Asymptotic behaviour of Green function

Let us consider the following boundary value problem:

$$(2.1) \quad \frac{\partial u}{\partial t} = \Delta u - Q(x)u, \quad x \in \mathbb{R}_n^+, \quad t > 0,$$

$$(2.2) \quad u(x, 0) = \Psi(x), \quad x \in \mathbb{R}_n^+, \quad \Psi(x) \in H_1$$

$$(2.3) \quad u(t, x)|_{x_n=0} = 0.$$

Let $G(t, x, s)$ be the Green function of this problem. The Green function for the problem (2.1)-(2.3) corresponding $Q(x) = 0$ is

$$(2.4) \quad G_0(t, x, s) = \left(2\sqrt{\pi t}\right)^{-n} \left[e^{-\frac{|x-s|^2}{4t}} - e^{-\frac{|x-s'|^2}{4t}} \right] I,$$

where $s = (s_1, s_2, s_3)$ and $s' = (s_1, s_2, -s_3)$, ($s_3 > 0$). We shall seek, using the method of E.E.Levi [4], the Green function of (1.1)-(1.3) in the form

$$(2.5) \quad G(t, x, s) = e^{-tQ(s)} G_0(t, x, s) + \int_0^t d\tau \int_{\mathbb{R}_n^+} G_0(t-\tau, x, \xi) e^{-(t-\tau)Q(\xi)} \varphi(\tau, \xi, s) d\xi,$$

where $\varphi(t, x, s)$ is a operator valued function to be determined. In order that $G(t, x, s)$ shall be a solution of (2.1) it is necessary that $\varphi(t, x, s)$ shall be a solution of the integral equation

$$(2.6) \quad \varphi(t, x, s) = K(t, x, s) + \int_0^t d\tau \int_{\mathbb{R}_n^+} K(t-\tau, x, \xi) \varphi(\tau, \xi, s) d\xi,$$

where

$$(2.7) \quad K(t, x, s) = [Q(s) - Q(x)] e^{-tQ(s)} G_0(t, x, s)$$

If we solve the equations by the method of successive approximation with the initial approximation $K_1(t, x, s) = K(t, x, s)$, then we obtain the iteration formula:

$$(2.8) \quad K_{n+1}(t, x, s) = \int_0^t d\tau \int_{\mathbb{R}_n^+} K(t-\tau, x, \xi) K_n(\tau, \xi, s) d\xi, \quad n = 1, 2, \dots$$

If the conditions (1^0) - (4^0) are satisfied then it can be shown that the series $\sum_{k=1}^{\infty} \|K_k(t, x, s)\|_1$ is uniformly convergent, $t > 0$, $x, s \in \mathbb{R}_n^+$. Therefore $\varphi(t, x, s) = \sum_{n=1}^{\infty} K_n(t, x, s)$ is a solution of the integral equation (2.6). Finding an estimate for $K_n(t, x, s)$ and $Q^\gamma(x) \varphi(t, x, s)$ we conclude our results with the following theorem.

Theorem 1. *If $Q(x)$ satisfies the conditions (1^0) - (4^0) , then the following asymptotic formula holds*

$$Q^\gamma(x) G(t, x, s) = Q^\gamma(x) e^{-tQ(s)} G_0(t, x, s) + O(1) t^{-\frac{n}{2} + \delta} e^{-\frac{|x-s|^2}{t}} \left[\|Q^{-k}(s)\|_1 + \|Q^\gamma(s) e^{-ctQ(s)}\|_1 \right],$$

where δ is a positive number.

§3. Asymptotic behaviour of $N_\beta(\lambda)$

Since the Green function of (2.1)-(2.2) is $G(t, x, s)$, then

$$(3.1) \quad u(t, x) = \int_{\mathbb{R}_n^+} G(t, x, s) \Psi(s) ds.$$

On the other hand, since $u(t, x) = e^{-tL}\psi(x)$, then we may write

$$(3.2) \quad e^{-tL}\Psi(x) = \int_{\mathbb{R}_n^+} G(t, x, s) \Psi(s) ds.$$

Therefore e^{-tL} is an integral operator with the kernel $G(t, x, s)$. Using the conditions (5⁰) and from the Theorem 1 we have

$$(3.3) \quad \int_{\mathbb{R}_n^+} Tr [Q^\gamma G(t, x, x)] dx \sim (2\sqrt{\pi t})^{-n} \int_{\mathbb{R}_n^+} Tr [Q^\gamma e^{-tQ(x)}] dx.$$

Considering (3.3) we obtain

$$\int_0^\infty Tr G(t, x, x) dx < \infty.$$

Since e^{-tL} is a positive operator we may write [1]

$$(3.4) \quad Tre^{-tL} = \int_{\mathbb{R}_n^+} Tr G(t, x, x) dx.$$

This implies that e^{-tL} is the trace operator in $H_1 = L_2(\mathbb{R}_n^+; H)$ and therefore the spectrum of the operator L is discrete: $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \leq \dots$, $\lim_{n \rightarrow \infty} \lambda_n = \infty$. It can be shown that $Q^\beta e^{-tL}$ is a trace operator in H_1 and that

$$(3.5) \quad Tr [Q^\beta e^{-tL}] = \int_{\mathbb{R}_n^+} Tr [Q^\beta G(t, x, x)] dx = \int_0^\infty e^{-\lambda t} N_\beta(\lambda) d\lambda$$

From (3.3) and (3.5) we have

$$(3.6) \quad \int_0^\infty e^{-\lambda t} N_\beta(\lambda) d\lambda \sim (2\sqrt{\pi t})^{-n} \int_{\mathbb{R}_n^+} Tr [Q^\beta e^{-tQ(x)}] dx.$$

As in [2] we write

$$(3.7) \quad (2\sqrt{\pi t})^{-n} \int_{\mathbb{R}_n^+} \alpha_j^\beta(x) e^{-t\alpha_j(x)} dx = \int_0^\infty e^{-\lambda t} d\rho_j(\lambda),$$

where $\rho_j(\lambda) = \frac{1}{2^n \pi^{\frac{n}{2}} \Gamma(\frac{n}{2}+1)} \sum_j \int_{\alpha_j(x) \leq \lambda} \alpha_j^\beta(x) [\lambda - \alpha_j(x)]^{n/2} dx$.

Using the inequality above we get

$$(3.8) \quad (2\sqrt{\pi t})^{-n} \int_{\mathbb{R}_n^+} \text{Tr} [Q^\beta e^{-tQ(x)}] dx = \int_0^\infty e^{-\lambda t} d\rho_\beta(\lambda).$$

Using (3.6) and (3.8) we obtain

$$(3.9) \quad \int_0^\infty e^{-\lambda t} N_\beta(\lambda) d\lambda = \int_0^\infty e^{-\lambda t} d\rho_\beta(\lambda) \text{ as } t \rightarrow 0.$$

Using the condition (6⁰) and application of a Tauberian theorem of Korenblum, B.I., (see [3]) to (3.9) we obtain that

$$(3.10) \quad N_\beta(\lambda) \sim \frac{1}{2^n \pi^{\frac{n}{2}} \Gamma(\frac{n}{2}+1)} \sum_j \int_{\alpha_j(x) \leq \lambda} \alpha_j^\beta(x) [\lambda - \alpha_j(x)]^{n/2} dx$$

as $\lambda \rightarrow \infty$.

Theorem2. *If $Q(x)$ satisfies the conditions (1⁰)-(6⁰), then the asymptotic formula (3.10) holds for the weighted trace $N_\beta(\lambda)$ of the operator L .*

References

- [1] A.A. Krillov, *Elementary Theory of Representations*, Springer Verlag, New York, 1976.
- [2] A.G. Kostjucenko, *Asymptotics of the spectral function of a self-adjoint elliptic operator*, Mathematics Summer School, Kiev, 1978.
- [3] B.I. Korenblum, *General Tauberian theorem for the rate of functions*, Dokl. Akad. Nauk. SSSR 88 (1953), 745-748.
- [4] S.D. Eidel'man, *Parabolic Systems*, North-Holland Publishing Company, Amsterdam 1969.
- [5] T. Kato, *Perturbation Theory for Linear Operators*, Springer Verlag, 1980.

Authors' address:

Fatih Taşçi and Mehmet Bayramoğlu
Yildiz Technical University, Mathematical Engineering Department,
Davutpasa Campus, 34010 Topkapi, Istanbul, Turkey.
E-mail addresses: tasci@yildiz.edu.tr, mbayram@yildiz.edu.tr