

Submanifolds in metallic Riemannian manifolds

Cristina E. Hretcanu and Adara M. Blaga

Abstract. The aim of our paper is to focus on some properties of submanifolds in Riemannian manifolds endowed with endomorphisms that generalize the Golden Riemannian structure, named metallic Riemannian structures. We focus on the properties of the structure induced on submanifolds, named by us Σ -metallic Riemannian structures, especially regarding the normality of this types of structure. Examples of structures induced on a sphere of codimension 1 by some metallic Riemannian structures defined on an Euclidean space are given.

M.S.C. 2010: 53B20, 53B25, 53C15.

Key words: Golden Riemannian structure; metallic Riemannian structure; normality of Riemannian structure; induced structure on submanifold.

1 Introduction

Metallic means family was introduced by Vera W. de Spinadel in ([10]) and it contains some generalizations of the Golden mean, such as *the Silver mean*, *the Bronze mean*, *the Copper mean*, *the Nickel mean* and many others.

A (p, q) -metallic number is the positive solution of the equation $x^2 - px - q = 0$ (for fixed positive integer values of p and q) and it has the form:

$$(1.1) \quad \sigma_{p,q} = \frac{p + \sqrt{p^2 + 4q}}{2}.$$

Some important members of the metallic mean family ([10]) are the following: the Golden mean $\phi = \frac{1+\sqrt{5}}{2}$ (for $p = q = 1$), the Silver mean $\sigma_{Ag} = \sigma_{2,1} = 1 + \sqrt{2}$ (for $q = 1$ and $p = 2$), the Bronze mean $\sigma_{Br} = \sigma_{3,1} = \frac{3+\sqrt{13}}{2}$ (for $q = 1$ and $p = 3$), the Subtle mean $\sigma_{4,1} = 2 + \sqrt{5} = \phi^3$ (for $p = 4$ and $q = 1$), the Copper mean $\sigma_{Cu} = \sigma_{1,2} = 2$ (for $p = 1$ and $q = 2$), the Nickel mean $\sigma_{Ni} = \sigma_{1,3} = \frac{1+\sqrt{13}}{2}$ (for $p = 1$ and $q = 3$) and so on.

Polynomial structures on manifolds were defined by S.I. Goldberg, K. Yano and N.C. Petridis in ([3] and [4]). C.E. Hretcanu and M. Crasmareanu defined some particular cases of polynomial structures, called *Golden structure* ([6], [7], [1], [2]) and some generalizations of this, called *metallic structure* ([8]).

In this paper, after recalling the basic notions of Golden and metallic Riemannian manifolds (in Section 2), we focus on the properties of Σ -metallic Riemannian structures induced on submanifolds by metallic Riemannian structure (in Section 3) and we give some examples of structures induced on submanifolds by metallic Riemannian structures (in Section 4). In the next section we find some properties of the normal Σ -metallic Riemannian structures.

2 Golden and metallic Riemannian structures

The *Golden structure* ([1]) is a polynomial structure on a manifold \overline{M} determined by a $(1,1)$ -tensor field J which satisfies:

$$(2.1) \quad J^2 = J + I,$$

where I is the identity operator on the Lie algebra $\mathcal{X}(\overline{M})$ of vector fields on \overline{M} .

A *metallic structure* ([8]) is a polynomial structure on a manifold \overline{M} determined by a $(1,1)$ -tensor field J which satisfies:

$$(2.2) \quad J^2 = pJ + qI,$$

where I is the identity operator on the Lie algebra $\mathcal{X}(\overline{M})$ of vector fields on \overline{M} and p, q are fixed positive integer numbers.

Moreover, if $(\overline{M}, \overline{g})$ is a Riemannian manifold endowed with a metallic structure J such that the Riemannian metric \overline{g} is J -compatible, i.e.:

$$(2.3) \quad \overline{g}(JX, Y) = \overline{g}(X, JY),$$

for any $X, Y \in \mathcal{X}(\overline{M})$, then (\overline{g}, J) is called *metallic Riemannian structure* and $(\overline{M}, \overline{g}, J)$ is a *metallic Riemannian manifold*.

In a metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$, from (2.2) and (2.3) we get:

$$(2.4) \quad \overline{g}(JX, JY) = p\overline{g}(X, JY) + q\overline{g}(X, Y),$$

for any $X, Y \in \mathcal{X}(\overline{M})$.

An almost product structure F on a Riemannian manifold $(\overline{M}, \overline{g})$ induces two metallic Riemannian structures on \overline{M} , given by ([8]):

$$(2.5) \quad J_1 = \frac{p}{2}I + \left(\frac{2\sigma_{p,q} - p}{2}\right)F, \quad J_2 = \frac{p}{2}I - \left(\frac{2\sigma_{p,q} - p}{2}\right)F.$$

Conversely, every metallic structure J on a Riemannian manifold $(\overline{M}, \overline{g})$ induces two almost product structures on this manifold ([8]):

$$(2.6) \quad F_{\pm} = \pm \left(\frac{2}{2\sigma_{p,q} - p} J - \frac{p}{2\sigma_{p,q} - p} I \right).$$

3 Σ -metallic Riemannian structures

In this section we assume that M is an n -dimensional submanifold isometrically immersed in the $(n+r)$ -dimensional metallic Riemannian manifold (\bar{M}, \bar{g}, J) , with $n, r \geq 1$ integer numbers.

We denote by $T_x(M)$ the tangent space of M in a point $x \in M$ and by $T_x^\perp(M)$ the normal space of M in x . Let i_* be the differential of the immersion $i : M \rightarrow \bar{M}$. The induced Riemannian metric g on M is given by $g(X, Y) = \bar{g}(i_*X, i_*Y)$, for any $X, Y \in \mathcal{X}(M)$. We fix a local orthonormal basis $\{N_1, \dots, N_r\}$ of the normal space $T_x^\perp(M)$. Hereafter we assume that the indices α, β, γ run over the range $\{1, \dots, r\}$.

The vector fields $J(i_*X)$ and $J(N_\alpha)$ can be decomposed into tangential and normal components:

$$(3.1) \quad (i) \ Ji_*X = i_*(PX) + \sum_{\alpha=1}^r u_\alpha(X)N_\alpha, \quad (ii) \ JN_\alpha = i_*(\xi_\alpha) + \sum_{\beta=1}^r a_{\alpha\beta}N_\beta,$$

for any $X \in T_x(M)$, where P is a $(1, 1)$ -tensor field on M , ξ_α are vector fields on M , u_α are 1-forms on M and $(a_{\alpha\beta})_r$ is an $r \times r$ matrix of smooth real functions on M .

In the rest of the paper we shall simply denote by X the vector field i_*X , for $X \in \mathcal{X}(M)$.

Theorem 3.1. ([8]) *The structure $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$, induced on a submanifold M by the metallic Riemannian structure (\bar{g}, J) on \bar{M} , satisfies the following equalities:*

$$(3.2) \quad P^2X = pPX + qX - \sum_{\alpha} u_\alpha(X)\xi_\alpha,$$

$$(3.3) \quad (i) \ u_\alpha(PX) = pu_\alpha(X) - \sum_{\beta} a_{\alpha\beta}u_\beta(X), \quad (ii) \ a_{\alpha\beta} = a_{\beta\alpha},$$

$$(3.4) \quad (i) \ u_\beta(\xi_\alpha) = q\delta_{\alpha\beta} + pa_{\alpha\beta} - \sum_{\gamma} a_{\alpha\gamma}a_{\gamma\beta}, \quad (ii) \ P\xi_\alpha = p\xi_\alpha - \sum_{\beta} a_{\alpha\beta}\xi_\beta,$$

$$(3.5) \quad (i) \ g(PX, Y) = g(X, PY), \quad (ii) \ u_\alpha(X) = g(X, \xi_\alpha)$$

for any $X, Y \in \mathcal{X}(M)$, where $\delta_{\alpha\beta}$ is the Kronecker delta and p, q are fixed positive integers.

Definition 3.1. A Σ -metallic Riemannian structure on a Riemannian manifold (M, g) is given by the data $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$, determined by a $(1, 1)$ -tensor field P on M , the vector fields ξ_α on M , the 1-forms u_α on M , the $r \times r$ matrix $(a_{\alpha\beta})_r$ of smooth real functions on M which verify the relations (3.2), (3.3), (3.4) and (3.5).

Remark 3.2. If $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$ is the induced structure on the submanifold M in the metallic Riemannian manifold (\bar{M}, \bar{g}, J) , then M is an invariant submanifold with respect to J (i.e. $J(M) \subset TM$) if and only if (M, g, P) is a metallic Riemannian manifold, whenever P is non-trivial ([8]).

Let $\bar{\nabla}$ and ∇ be the Levi-Civita connections on (\bar{M}, \bar{g}) and (M, g) , respectively. The Gauss and Weingarten formulas are:

$$(3.6) \quad (i) \bar{\nabla}_X Y = \nabla_X Y + \sum_{\alpha=1}^r h_\alpha(X, Y) N_\alpha, \quad (ii) \bar{\nabla}_X N_\alpha = -A_\alpha X + \nabla_X^\perp N_\alpha,$$

respectively, where $A_\alpha =: A_{N_\alpha}$ is the shape operator and $h(X, Y) = \sum_{\alpha=1}^r h_\alpha(X, Y) N_\alpha$ is the second fundamental form. Also,

$$(3.7) \quad h_\alpha(X, Y) = g(h(X, Y), N_\alpha) = g(A_\alpha X, Y),$$

for any $X, Y \in \mathcal{X}(M)$.

Remark 3.3. The normal connection $\nabla_X^\perp N_\alpha$ has the decomposition:

$$(3.8) \quad \nabla_X^\perp N_\alpha = \sum_{\beta=1}^r l_{\alpha\beta}(X) N_\beta,$$

for any $X \in \mathcal{X}(M)$, $\alpha \in \{1, \dots, r\}$, and

$$(3.9) \quad l_{\alpha\beta} = -l_{\beta\alpha},$$

for any $\alpha, \beta \in \{1, \dots, r\}$.

Proof. From $\bar{g}(\nabla_X^\perp N_\alpha, N_\beta) + \bar{g}(N_\alpha, \nabla_X^\perp N_\beta) = 0$ we get:

$$\bar{g}\left(\sum_{\gamma} l_{\alpha\gamma}(X) N_\gamma, N_\beta\right) + \bar{g}\left(N_\alpha, \sum_{\gamma} l_{\beta\gamma}(X) N_\gamma\right) = 0,$$

for any $X \in \mathcal{X}(M)$ and $\gamma \in \{1, \dots, r\}$. Thus we obtain (3.9). \square

Using an analogy of a locally product manifold ([9]), we can define *locally metallic Riemannian manifold*, as follows:

Definition 3.4. If (\bar{M}, \bar{g}, J) is a metallic Riemannian manifold and J is parallel with respect to the Levi-Civita connection $\bar{\nabla}$ on \bar{M} (i.e. $\bar{\nabla} J = 0$), then we call (\bar{M}, \bar{g}, J) a *locally metallic Riemannian manifold*.

As in the case of submanifolds in Riemannian manifolds with Golden structure ([6]), it is easy to verify the following relations regarding the covariant derivatives of components of the $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$ -structure, induced on M by the metallic Riemannian structure (\bar{g}, J) :

Theorem 3.2. *If M is an n -dimensional submanifold of codimension r in a locally metallic Riemannian manifold (\bar{M}, \bar{g}, J) , then the $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$ -structure, induced on M by the metallic Riemannian structure (\bar{g}, J) , has the following properties:*

$$(3.10) \quad (\nabla_X P)(Y) = \sum_{\alpha=1}^r h_\alpha(X, Y) \xi_\alpha + \sum_{\alpha=1}^r u_\alpha(Y) A_\alpha X,$$

$$(3.11) \quad (\nabla_X u_\alpha)(Y) = -h_\alpha(X, PY) + \sum_{\beta=1}^r [l_{\alpha\beta}(X)u_\beta(Y) + a_{\beta\alpha}h_\beta(X, Y)],$$

$$(3.12) \quad \nabla_X \xi_\alpha = -P(A_\alpha X) + \sum_{\beta=1}^r a_{\alpha\beta}A_\beta X + \sum_{\beta=1}^r l_{\alpha\beta}(X)\xi_\beta,$$

$$(3.13) \quad X(a_{\alpha\beta}) = -u_\alpha(A_\beta X) - u_\beta(A_\alpha X) + \sum_{\gamma=1}^r [a_{\gamma\beta}l_{\alpha\gamma}(X) + a_{\alpha\gamma}l_{\beta\gamma}(X)],$$

for any $X, Y \in \mathcal{X}(M)$.

Proof. Using the Gauss and Weingarten formulae in $(\bar{\nabla}_X J)Y = 0$, for any $X, Y \in \mathcal{X}(M)$ and identifying the tangential components (and the normal components, respectively), we obtain the equalities (3.10) and (3.11). Moreover, from

$$\bar{\nabla}_X(JN_\alpha) = \nabla_X \xi_\alpha - \sum_{\beta=1}^r a_{\alpha\beta}A_\beta X + \sum_{\beta=1}^r [X(a_{\alpha\beta}) + h_\beta(X, \xi_\alpha) + \sum_{\gamma=1}^r a_{\alpha\gamma}l_{\gamma\beta}(X)]N_\beta$$

and

$$J(\bar{\nabla}_X N_\alpha) = -P(A_\alpha X) + \sum_{\beta=1}^r l_{\alpha\beta}(X)\xi_\beta - \sum_{\beta=1}^r [u_\beta(A_\alpha X) - \sum_{\gamma=1}^r a_{\gamma\beta}l_{\alpha\gamma}(X)]N_\beta,$$

for any $X \in \mathcal{X}(M)$ and using $(\bar{\nabla}_X J)N_\alpha = 0$ we obtain (3.12) and (3.13) by identifying the tangential and the normal components, respectively. \square

Remark 3.5. If M is an invariant submanifold in the locally metallic Riemannian manifold (\bar{M}, \bar{g}, J) , then P is parallel with respect to the Levi-Civita connection ∇ on M , where P is the $(1, 1)$ -tensor field on the Riemannian manifold (M, g) defined in (3.1)(i).

4 Examples of structures induced on submanifolds by metallic Riemannian structures

We assume that the ambient space is E^{2a+b} ($a, b \geq 1$ integer numbers) and for any point of E^{2a+b} we have its coordinates:

$$(x^1, \dots, x^a, y^1, \dots, y^a, z^1, \dots, z^b) := (x^i, y^i, z^j),$$

where $i \in \{1, \dots, a\}$ and $j \in \{1, \dots, b\}$. The tangent space $T_x(E^{2a+b})$ is isomorphic with E^{2a+b} . For $\lambda \in \{-1, 1\}$ we can construct two metallic structures on E^{2a+b} , $J_\lambda : E^{2a+b} \rightarrow E^{2a+b}$, given by:

$$(4.1) \quad J_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{p}{2} \frac{\partial}{\partial x^i} + \frac{\lambda\sqrt{\Delta}}{2} \frac{\partial}{\partial y^i}, \frac{p}{2} \frac{\partial}{\partial y^i} + \frac{\lambda\sqrt{\Delta}}{2} \frac{\partial}{\partial x^i}, \left(\frac{p}{2} + \frac{\lambda\varepsilon_j\sqrt{\Delta}}{2} \right) \frac{\partial}{\partial z^j} \right),$$

where $\varepsilon_j \in \{-1, 1\}$, $i \in \{1, \dots, a\}$, $j \in \{1, \dots, b\}$ and $\Delta = p^2 + 4q$.

We easily find that $J_\lambda^2 = pJ_\lambda + qI$ and $(J_\lambda, \langle \cdot, \cdot \rangle)$ is a metallic Riemannian structure on E^{2a+b} .

Thus, we obtain two metallic structures: J_{+1} (for $\lambda = 1$) and J_{-1} (for $\lambda = -1$). Moreover, for $\lambda\varepsilon_j = 1$, we obtain the following form of J_λ :

$$(4.2) \quad J_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{p}{2} \frac{\partial}{\partial x^i} + \lambda \frac{\sqrt{\Delta}}{2} \frac{\partial}{\partial y^i}, \frac{p}{2} \frac{\partial}{\partial y^i} + \lambda \frac{\sqrt{\Delta}}{2} \frac{\partial}{\partial x^i}, \sigma_{p,q} \frac{\partial}{\partial z^j} \right),$$

and for $\lambda\varepsilon_j = -1$, we obtain:

$$(4.3) \quad J_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{p}{2} \frac{\partial}{\partial x^i} + \lambda \frac{\sqrt{\Delta}}{2} \frac{\partial}{\partial y^i}, \frac{p}{2} \frac{\partial}{\partial y^i} + \lambda \frac{\sqrt{\Delta}}{2} \frac{\partial}{\partial x^i}, \bar{\sigma}_{p,q} \frac{\partial}{\partial z^j} \right),$$

where $\bar{\sigma}_{p,q} = p - \sigma$. Now, we can construct a Σ -metallic Riemannian structure on the sphere $S^{2a+b-1}(R) \hookrightarrow E^{2a+b}$. The equation of sphere $S^{2a+b-1}(R)$ is:

$$(4.4) \quad \sum_{i=1}^a (x^i)^2 + \sum_{i=1}^a (y^i)^2 + \sum_{j=1}^b (z^j)^2 = R^2,$$

where R is its radius and $(x^1, \dots, x^a, y^1, \dots, y^a, z^1, \dots, z^b) := (x^i, y^i, z^j)$ are the coordinates of any point of $S^{2a+b-1}(R)$. We denote by:

$$(4.5) \quad \sum_{i=1}^a (x^i)^2 = r_1^2, \quad \sum_{i=1}^a (y^i)^2 = r_2^2, \quad \sum_{j=1}^b (z^j)^2 = r_3^2$$

and we obtain $r_1^2 + r_2^2 + r_3^2 = R^2$.

An unit normal vector field N on the sphere $S^{2a+b-1}(R)$ is given by:

$$(4.6) \quad N = \frac{1}{R} (x^1, \dots, x^a, y^1, \dots, y^a, z^1, \dots, z^b).$$

Thus:

$$(4.7) \quad J_\lambda N = \frac{1}{R} \left(\frac{p}{2} x^i + \lambda \frac{\sqrt{\Delta}}{2} y^i, \frac{p}{2} y^i + \lambda \frac{\sqrt{\Delta}}{2} x^i, \sigma_{p,q} z^j \right),$$

where $\Delta = p^2 + 4q$ and $\lambda \in \{-1, 1\}$.

For a vector field X on $S^{2a+b-1}(R)$ we use the following notation:

$$X = (X^1, \dots, X^a, Y^1, \dots, Y^a, Z^1, \dots, Z^b) := (X^i, Y^i, Z^j)$$

and from $\langle X, N \rangle = 0$ we obtain:

$$(4.8) \quad \sum_{i=1}^a (x^i X^i + y^i Y^i) + \sum_{j=1}^b z^j Z^j = 0.$$

If we decompose $J_\lambda N$ into tangential and normal components at the sphere $S^{2a+b-1}(R)$, for $\mathcal{A} = \langle J_\lambda N, N \rangle$, we obtain:

$$(4.9) \quad J_\lambda N = \xi + \mathcal{A}N.$$

In the followings we consider $\lambda \varepsilon_j = 1$. Thus, we obtain:

$$(4.10) \quad \mathcal{A} = \frac{1}{R^2} \left[\frac{p}{2}(r_1^2 + r_2^2) + \lambda \sqrt{\Delta} \sum_{i=1}^a x^i y^i + \sigma_{p,q} r_3^2 \right].$$

The tangential component of $J_\lambda N$ at the sphere $S^{2a+b-1}(R)$ has the form:

$$(4.11) \quad \xi = \frac{1}{R} \left(\left(\frac{p}{2} - \mathcal{A} \right) x^i + \lambda \frac{\sqrt{\Delta}}{2} y^i, \left(\frac{p}{2} - \mathcal{A} \right) y^i + \lambda \frac{\sqrt{\Delta}}{2} x^i, (\sigma_{p,q} - \mathcal{A}) z^j \right).$$

From $u(X) = \langle X, \xi \rangle$, we obtain:

$$(4.12) \quad u(X) = \lambda \frac{\sqrt{\Delta}}{2R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right].$$

The decomposition of JX into tangential and normal components at the sphere $S^{2a+b-1}(R)$ (where X is a vector field on $S^{2a+b-1}(R)$), is given by:

$$(4.13) \quad J_\lambda X = PX + u(X)N.$$

Thus, we obtain:

$$(4.14) \quad PX = \left(\frac{p}{2} X^i + \lambda \frac{\sqrt{\Delta}}{2} Y^i - \frac{u(X)}{R} x^i, \frac{p}{2} Y^i + \lambda \frac{\sqrt{\Delta}}{2} X^i - \frac{u(X)}{R} y^i, \sigma_{p,q} Z^j - \frac{u(X)}{R} z^j \right),$$

for any tangent vector $X := (X^i, Y^i, Z^j)$ at the sphere $S^{2a+b-1}(R)$ in a point (x^i, y^i, z^j) , for $i \in \{1, \dots, a\}$ and $j \in \{1, \dots, b\}$.

Therefore, from the relations (4.10), (4.11), (4.12) and (4.14) we obtain the Σ -structure induced on the sphere $S^{2a+b-1}(R)$ of codimension 1 in the Euclidean space E^{2a+b} by J_λ from E^{2a+b} , given by $(P, \langle \cdot, \cdot \rangle, \xi, u, \mathcal{A})$.

In particular, for $\lambda \varepsilon_j = 1$ (where $\lambda \in \{-1, 1\}$ and $\varepsilon_j \in \{-1, 1\}$), we obtain:

1. for $p = q = 1$, two Golden structures, given by:

$$(4.15) \quad \Phi_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{1}{2} \frac{\partial}{\partial x^i} + \lambda \frac{\sqrt{5}}{2} \frac{\partial}{\partial y^i}, \frac{1}{2} \frac{\partial}{\partial y^i} + \lambda \frac{\sqrt{5}}{2} \frac{\partial}{\partial x^i}, \phi \frac{\partial}{\partial z^j} \right)$$

2. for $p = 2, q = 1$, two Silver structures, given by:

$$(4.16) \quad Ag_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{\partial}{\partial x^i} + \lambda \sqrt{2} \frac{\partial}{\partial y^i}, \frac{\partial}{\partial y^i} + \lambda \sqrt{2} \frac{\partial}{\partial x^i}, \sigma_{Ag} \frac{\partial}{\partial z^j} \right)$$

3. for $p = 3, q = 1$, two Bronze structures, given by:

$$(4.17) \quad Br_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{3}{2} \frac{\partial}{\partial x^i} + \lambda \frac{\sqrt{13}}{2} \frac{\partial}{\partial y^i}, \frac{3}{2} \frac{\partial}{\partial y^i} + \lambda \frac{\sqrt{13}}{2} \frac{\partial}{\partial x^i}, \sigma_{Br} \frac{\partial}{\partial z^j} \right)$$

4. for $p = 1, q = 2$, two Copper structures, given by:

$$(4.18) \quad Cu_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{1}{2} \frac{\partial}{\partial x^i} + \lambda \frac{3}{2} \frac{\partial}{\partial y^i}, \frac{1}{2} \frac{\partial}{\partial y^i} + \lambda \frac{3}{2} \frac{\partial}{\partial x^i}, \sigma_{Cu} \frac{\partial}{\partial z^j} \right)$$

5. for $p = 1, q = 3$, two Nickel structures, given by:

$$(4.19) \quad Ni_\lambda \left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i}, \frac{\partial}{\partial z^j} \right) = \left(\frac{1}{2} \frac{\partial}{\partial x^i} + \lambda \frac{\sqrt{13}}{2} \frac{\partial}{\partial y^i}, \frac{1}{2} \frac{\partial}{\partial y^i} + \lambda \frac{\sqrt{13}}{2} \frac{\partial}{\partial x^i}, \sigma_{Ni} \frac{\partial}{\partial z^j} \right).$$

The induced structures on the sphere $S^{2a+b-1}(R)$ of codimension 1 in the Euclidean space E^{2a+b} are as follows:

1. for $p = q = 1$, we obtain the Σ -Golden structures, given by:

$$\begin{aligned} \mathcal{A} &= \frac{1}{R^2} \left[\frac{1}{2}(r_1^2 + r_2^2) + \lambda\sqrt{5} \sum_{i=1}^a x^i y^i + \phi r_3^2 \right] \\ \xi &= \frac{1}{R} \left(\left(\frac{1}{2} - \mathcal{A} \right) x^i + \lambda \frac{\sqrt{5}}{2} y^i, \left(\frac{1}{2} - \mathcal{A} \right) y^i + \lambda \frac{\sqrt{5}}{2} x^i, \left(\frac{1}{2} - \mathcal{A} + \frac{\sqrt{5}}{2} \right) z^j \right) \\ u(X) &= \lambda \frac{\sqrt{5}}{2R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right] \\ PX &= \left(\frac{1}{2} X^i + \lambda \frac{\sqrt{5}}{2} Y^i - \frac{u(X)}{R} x^i, \frac{1}{2} Y^i + \lambda \frac{\sqrt{5}}{2} X^i - \frac{u(X)}{R} y^i, \phi Z^j - \frac{u(X)}{R} z^j \right), \end{aligned}$$

induced by the structure defined in (4.15);

2. for $p = 2, q = 1$, we obtain the Σ -Silver structures, given by:

$$\begin{aligned} \mathcal{A} &= \frac{1}{R^2} \left[(r_1^2 + r_2^2) + 2\lambda\sqrt{2} \sum_{i=1}^a x^i y^i + \sigma_{Ag} r_3^2 \right] \\ \xi &= \frac{1}{R} \left((1 - \mathcal{A}) x^i + \lambda\sqrt{2} y^i, (1 - \mathcal{A}) y^i + \lambda\sqrt{2} x^i, (1 - \mathcal{A} + \sqrt{2}) z^j \right) \\ u(X) &= \lambda \frac{\sqrt{2}}{R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right] \\ PX &= \left(X^i + \lambda\sqrt{2} Y^i - \frac{u(X)}{R} x^i, Y^i + \lambda\sqrt{2} X^i - \frac{u(X)}{R} y^i, \sigma_{Ag} Z^j - \frac{u(X)}{R} z^j \right), \end{aligned}$$

induced by the structure defined in (4.16);

3. for $p = 3, q = 1$, we obtain the Σ -Bronze structures, given by:

$$\begin{aligned} \mathcal{A} &= \frac{1}{R^2} \left[\frac{3}{2}(r_1^2 + r_2^2) + \lambda\sqrt{13} \sum_{i=1}^a x^i y^i + \sigma_{Br} r_3^2 \right] \\ \xi &= \frac{1}{R} \left(\left(\frac{3}{2} - \mathcal{A} \right) x^i + \lambda \frac{\sqrt{13}}{2} y^i, \left(\frac{3}{2} - \mathcal{A} \right) y^i + \lambda \frac{\sqrt{13}}{2} x^i, \left(\frac{3}{2} - \mathcal{A} + \frac{\sqrt{13}}{2} \right) z^j \right) \\ u(X) &= \lambda \frac{\sqrt{13}}{2R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right] \\ PX &= \left(\frac{3}{2} X^i + \lambda \frac{\sqrt{13}}{2} Y^i - \frac{u(X)}{R} x^i, \frac{3}{2} Y^i + \lambda \frac{\sqrt{13}}{2} X^i - \frac{u(X)}{R} y^i, \sigma_{Br} Z^j - \frac{u(X)}{R} z^j \right) \end{aligned}$$

induced by the structure defined in (4.17);

4. for $p = 1, q = 2$, we obtain the Σ -Copper structures, given by:

$$\begin{aligned} \mathcal{A} &= \frac{1}{R^2} \left[\frac{1}{2}(r_1^2 + r_2^2) + 3\lambda \sum_{i=1}^a x^i y^i + \sigma_{Cu} r_3^2 \right] \\ \xi &= \frac{1}{R} \left(\left(\frac{1}{2} - \mathcal{A} \right) x^i + \lambda \frac{3}{2} y^i, \left(\frac{1}{2} - \mathcal{A} \right) y^i + \lambda \frac{3}{2} x^i, \left(\frac{1}{2} - \mathcal{A} + \frac{3}{2} \right) z^j \right) \\ u(X) &= \lambda \frac{3}{2R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right] \\ PX &= \left(\frac{1}{2} X^i + \lambda \frac{3}{2} Y^i - \frac{u(X)}{R} x^i, \frac{1}{2} Y^i + \lambda \frac{3}{2} X^i - \frac{u(X)}{R} y^i, \sigma_{Cu} Z^j - \frac{u(X)}{R} z^j \right) \end{aligned}$$

induced by the structure defined in (4.18);

5. for $p = 1, q = 3$, we obtain the Σ -Nickel structures, given by:

$$\begin{aligned} \mathcal{A} &= \frac{1}{R^2} \left[\frac{1}{2}(r_1^2 + r_2^2) + \lambda\sqrt{13} \sum_{i=1}^a x^i y^i + \sigma_{Ni} r_3^2 \right] \\ \xi &= \frac{1}{R} \left(\left(\frac{1}{2} - \mathcal{A} \right) x^i + \lambda \frac{\sqrt{13}}{2} y^i, \left(\frac{1}{2} - \mathcal{A} \right) y^i + \lambda \frac{\sqrt{13}}{2} x^i, \left(\frac{1}{2} - \mathcal{A} + \frac{\sqrt{13}}{2} \right) z^j \right) \\ u(X) &= \lambda \frac{\sqrt{13}}{2R} \left[\sum_{i=1}^a (y^i X^i + x^i Y^i) + \lambda \sum_{j=1}^b z^j Z^j \right] \\ PX &= \left(\frac{1}{2} X^i + \lambda \frac{\sqrt{13}}{2} Y^i - \frac{u(X)}{R} x^i, \frac{1}{2} Y^i + \lambda \frac{\sqrt{13}}{2} X^i - \frac{u(X)}{R} y^i, \sigma_{Ni} Z^j - \frac{u(X)}{R} z^j \right), \end{aligned}$$

induced by the structure defined in (4.19).

In conclusion, we can obtain Σ -metallic structures $(P, \langle \cdot, \cdot \rangle, \xi, u, \mathcal{A})$, induced on the sphere $S^{2a+b-1}(R)$ of codimension 1 in the Euclidean space E^{2a+b} by the metallic structure J_λ from E^{2a+b} , for various positive integer values of p and q .

5 On the normality of Σ -metallic Riemannian structures

Let N_J be the Nijenhuis torsion tensor field of J , defined by:

$$(5.1) \quad N_J(X, Y) := [JX, JY] + J^2[X, Y] - J[JX, Y] - J[X, JY],$$

for any $X, Y \in \mathcal{X}(\overline{M})$. It is known that the Nijenhuis tensor of J verifies ([11]):

$$(5.2) \quad N_J(X, Y) = (\overline{\nabla}_{JX}J)(Y) - (\overline{\nabla}_{JY}J)(X) - J[(\overline{\nabla}_XJ)(Y) - (\overline{\nabla}_YJ)(X)],$$

for any $X, Y \in \mathcal{X}(\overline{M})$.

Thus, we remark that if the metallic structure J on \overline{M} is parallel with respect to $\overline{\nabla}$, then $N_J = 0$.

We shall define a normal Σ -metallic Riemannian structure on the submanifold M in the metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$.

Definition 5.1. A Σ -metallic Riemannian structure defined on a submanifold (M, g) of codimension r in a Riemannian manifold $(\overline{M}, \overline{g})$ is said to be normal if:

$$(5.3) \quad N_P = 2 \sum_{\alpha=1}^r du_\alpha \otimes \xi_\alpha.$$

In the following considerations, let M be a submanifold in the locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$ and let ∇ be the Levi-Civita connection determined by the induced metric g on M .

We denote by $B_\alpha =: PA_\alpha - A_\alpha P$ and remark that $g(B_\alpha X, Y) = -g(B_\alpha Y, X)$, for any $X, Y \in \mathcal{X}(M)$.

Theorem 5.1. *If M is a submanifold of codimension r in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$ and $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$ is the structure induced on the Riemannian manifold (M, g) , then the Nijenhuis torsion tensor field of the $(1, 1)$ -tensor field P and the 1-forms u_α verify the equalities:*

$$(5.4) \quad N_P(X, Y) = \sum_{\alpha=1}^r [g(X, \xi_\alpha)B_\alpha Y - g(Y, \xi_\alpha)B_\alpha X - g(B_\alpha X, Y)\xi_\alpha],$$

$$(5.5) \quad 2du_\alpha(X, Y) = -g(B_\alpha X, Y) + \sum_{\beta=1}^r [l_{\alpha\beta}(X)g(Y, \xi_\beta) - l_{\alpha\beta}(Y)g(X, \xi_\beta)],$$

for any $X, Y \in \mathcal{X}(M)$, where $l_{\alpha\beta}$ are the coefficients of the normal connection ∇^\perp in the normal bundle $T^\perp(M)$.

Proof. By using (3.10) and (5.2), we get:

$$N_P(X, Y) = \sum_{\alpha=1}^r [g(A_\alpha PX, Y) - g(A_\alpha PY, X)]\xi_\alpha +$$

$$+ \sum_{\alpha=1}^r [g(Y, \xi_\alpha)A_\alpha PX - g(X, \xi_\alpha)A_\alpha PY] - \sum_{\alpha=1}^r [g(Y, \xi_\alpha)P(A_\alpha X) - g(X, \xi_\alpha)P(A_\alpha Y)]$$

and from $g(A_\alpha PY, X) = g(P(A_\alpha X), Y)$ we obtain that the Nijenhuis torsion tensor field of the $(1, 1)$ -tensor field P from the Σ -metallic Riemannian structure has the form:

$$N_P(X, Y) = - \sum_{\alpha=1}^r g((PA_\alpha - A_\alpha P)(X), Y)\xi_\alpha - \\ - \sum_{\alpha=1}^r g(Y, \xi_\alpha)(PA_\alpha - A_\alpha P)(X) + \sum_{\alpha=1}^r g(X, \xi_\alpha)(PA_\alpha - A_\alpha P)(Y),$$

for any $X, Y \in \mathcal{X}(M)$ which implies (5.4).

From $2du_\alpha(X, Y) = X(u_\alpha(Y)) - Y(u_\alpha(X)) - u_\alpha([X, Y])$ we have:

$$2du_\alpha(X, Y) = (\nabla_X u_\alpha)(Y) - (\nabla_Y u_\alpha)(X),$$

for any $X, Y \in \mathcal{X}(M)$ and using (3.11) we obtain:

$$2du_\alpha(X, Y) = g(A_\alpha Y, PX) - g(A_\alpha X, PY) + \\ + \sum_{\beta=1}^r [g(A_\beta X, Y) - g(A_\beta Y, X)]a_{\alpha\beta} + \sum_{\beta=1}^r [g(Y, \xi_\beta)l_{\alpha\beta}(X) - g(X, \xi_\beta)l_{\alpha\beta}(Y)].$$

From $g(A_\beta X, Y) = g(A_\beta Y, X)$ we have:

$$g(A_\alpha X, PY) - g(A_\alpha Y, PX) = g((PA_\alpha - A_\alpha P)(X), Y) = g(B_\alpha X, Y)$$

and we obtain that the 1-forms u_α of the Σ -metallic Riemannian structure induced on M verify the equality (5.5). \square

Remark 5.2. Let M be a submanifold of codimension r in a locally metallic Riemannian manifold (\bar{M}, \bar{g}, J) and let $\Sigma = (P, g, u_\alpha, \xi_\alpha, (a_{\alpha\beta})_r)$ be the structure induced on the Riemannian manifold (M, g) . If the $(1, 1)$ -tensor field P of the Σ -metallic Riemannian structure on M commutes with the Weingarten operators A_α , for any $\alpha \in \{1, \dots, r\}$ (i.e. $B_\alpha = 0$), then the Nijenhuis torsion tensor field of P vanishes on M (i.e. $N_P = 0$).

From (5.4) and (5.5) we obtain:

Theorem 5.2. *If M is a submanifold of codimension r in the locally metallic Riemannian manifold (\bar{M}, \bar{g}, J) , then:*

$$(5.6) \quad N_P(X, Y) - 2 \sum_{\alpha=1}^r du_\alpha(X, Y)\xi_\alpha = \sum_{\alpha=1}^r [g(X, \xi_\alpha)B_\alpha Y - g(Y, \xi_\alpha)B_\alpha X] - \\ - \sum_{\alpha=1}^r \sum_{\beta=1}^r [l_{\alpha\beta}(X)g(Y, \xi_\beta) - l_{\alpha\beta}(Y)g(X, \xi_\beta)]\xi_\alpha,$$

for any $X, Y \in \mathcal{X}(M)$, where $l_{\alpha\beta}$ are the coefficients of the normal connection in the normal bundle $T^\perp(M)$.

Remark 5.3. If the Σ -metallic Riemannian structure induced on M is normal and the normal connection ∇^\perp of M vanishes identically (i.e. $l_{\alpha\beta} = 0$), then we obtain:

$$(5.7) \quad \sum_{\alpha=1}^r g(X, \xi_\alpha)(PA_\alpha - A_\alpha P)(Y) = \sum_{\alpha=1}^r g(Y, \xi_\alpha)(PA_\alpha - A_\alpha P)(X)$$

and this equality does not depend on the choice of a basis in the normal space $T_x^\perp(M)$, for any $x \in M$.

Theorem 5.3. *Let M be a submanifold of codimension r in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$. If the normal connection ∇^\perp vanishes identically on the normal bundle $T^\perp(M)$ and the $(1, 1)$ -tensor field P of the Σ -metallic Riemannian structure induced on M commutes with every Weingarten operator A_α , then the induced Σ -metallic Riemannian structure on M is normal.*

Remark 5.4. The matrix $\mathcal{A} := (a_{\alpha\beta})_r$ of the structure Σ induced on an invariant submanifold M by the metallic structure J from the Riemannian manifold $(\overline{M}, \overline{g})$ is a metallic matrix, that is a matrix which verifies:

$$\mathcal{A}^2 = p\mathcal{A} + qI_r,$$

where I_r is the identically matrix of order r ([8]).

If $\mathcal{A} := (a_{\alpha\beta})_r$ is a metallic matrix, then $\sum_{\gamma=1}^r a_{\alpha\gamma}a_{\gamma\beta} = pa_{\alpha\beta} + q\delta_{\alpha\beta}$ and from (3.4)(i) we obtain $u_\beta(\xi_\alpha) = 0$, which implies that $P^2\xi_\alpha = pP\xi_\alpha + q\xi_\alpha$ and $J\xi_\alpha = P\xi_\alpha$, for any $\alpha \in \{1, \dots, r\}$.

Theorem 5.4. *Let M be a submanifold of codimension $r \geq 2$ in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$. If the normal connection ∇^\perp vanishes identically on the normal bundle $T^\perp(M)$ and M is a non-invariant submanifold with respect to the metallic structure J , then the vector fields $\{\xi_1, \dots, \xi_r\}$ are linearly independent.*

Proof. From (3.4)(i) we have:

$$g(\xi_\alpha, \xi_\beta) = q\delta_{\alpha\beta} + pa_{\alpha\beta} - \sum_{\gamma=1}^r a_{\alpha\gamma}a_{\gamma\beta}.$$

Let x_1, \dots, x_r be real numbers with the property $x_1\xi_1 + \dots + x_r\xi_r = 0$ in any point $x \in M$. Applying $g(\xi_\alpha, \cdot)$ in the last equality ($\alpha \in \{1, \dots, r\}$), we obtain a linear system, given by:

$$(5.8) \quad \sum_{i=1}^r x_i \Gamma_{ij} = 0$$

for any $j \in \{1, \dots, r\}$, where $\Gamma_{ii} = q + pa_{ii} - \sum_{\gamma} a_{i\gamma}^2$ and $\Gamma_{ij} = pa_{ij} - \sum_{\gamma} a_{i\gamma}a_{\gamma j}$ for $i \neq j$ and $i, j \in \{1, \dots, r\}$.

The determinant of the matrix of the linear system (5.8) is the determinant of the matrix $\mathcal{B} = qI_r + p\mathcal{A} - \mathcal{A}^2$, where $\mathcal{A} = (a_{\alpha\beta})_r$. Using the last remark, we observe that the determinant of the matrix \mathcal{B} is not zero when M is a non-invariant submanifold with respect to the metallic structure J . Therefore, the linear system of equations (5.8) has the unique solution $x_1 = \dots = x_r = 0$. Thus, the vector fields $\{\xi_1, \dots, \xi_r\}$ are linearly independent. \square

Theorem 5.5. *Let M be a submanifold of codimension $r \geq 1$ in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$. If the normal connection ∇^\perp vanishes identically on the normal bundle $T^\perp(M)$ and M is a non-invariant submanifold with respect to the metallic structure J , then the induced Σ -metallic Riemannian structure on M is normal if and only if the $(1,1)$ -tensor field P commutes with every Weingarten operator A_α , for any $\alpha \in \{1, \dots, r\}$.*

Proof. We assume that the induced Σ -metallic Riemannian structure on M is normal and we prove that the $(1,1)$ -tensor field P commutes with every Weingarten operator A_α (i.e. $PA_\alpha = A_\alpha P$), for any $\alpha \in \{1, \dots, r\}$.

If the normal connection vanishes identically (thus $l_{\alpha\beta} = 0$), then:

$$(5.9) \quad \sum_{\alpha=1}^r g(X, \xi_\alpha) B_\alpha Y = \sum_{\alpha=1}^r g(Y, \xi_\alpha) B_\alpha X.$$

Multiplying the equality (5.9) by $Z \in \mathcal{X}(M)$, we obtain:

$$(5.10) \quad \sum_{\alpha=1}^r g(X, \xi_\alpha) g(B_\alpha Y, Z) = \sum_{\alpha=1}^r g(Y, \xi_\alpha) g(B_\alpha X, Z),$$

for any $X, Y, Z \in \mathcal{X}(M)$.

Inverting Y by Z in the last equality and using that B_α is g -skew symmetric (i.e. $g(B_\alpha Y, Z) + g(B_\alpha Z, Y) = 0$), by summing these relations we obtain:

$$\sum_{\alpha=1}^r g(Y, \xi_\alpha) g(B_\alpha X, Z) + \sum_{\alpha=1}^r g(Z, \xi_\alpha) g(B_\alpha X, Y) = 0.$$

Therefore:

$$(5.11) \quad \sum_{\alpha=1}^r g(Y, \xi_\alpha) B_\alpha X + \sum_{\alpha=1}^r g(B_\alpha X, Y) \xi_\alpha = 0.$$

Inverting X by Y in the last equality and summing these relations we obtain:

$$\sum_{\alpha=1}^r g(Y, \xi_\alpha) B_\alpha X + \sum_{\alpha=1}^r g(X, \xi_\alpha) B_\alpha Y = 0$$

and multiplying this equality by $Z \in \mathcal{X}(M)$ we get:

$$(5.12) \quad \sum_{\alpha=1}^r g(Y, \xi_\alpha) g(B_\alpha X, Z) + \sum_{\alpha=1}^r g(X, \xi_\alpha) g(B_\alpha Y, Z) = 0.$$

Using (5.10) and (5.12) it follows that:

$$(5.13) \quad \sum_{\alpha=1}^r g(Y, \xi_\alpha) g(B_\alpha X, Z) = 0,$$

for any $X, Y, Z \in \mathcal{X}(M)$.

If M is non-invariant submanifold of codimension $r \geq 2$ in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$ and the normal connection ∇^\perp vanishes identically on the normal bundle $T^\perp(M)$, from the Theorem (5.4) we obtain that the vector fields ξ_1, \dots, ξ_r are linearly independent in any point $x \in M$. Due to these r linearly independent vector fields on M we have $r \leq n$. It follows that there exists a vector field $Y \in \mathcal{X}(M)$ which is orthogonal on the space spanned by $\{\xi_1, \dots, \xi_r\} - \{\xi_\alpha\}$ and $g(Y, \xi_\alpha) \neq 0$. Hence, we obtain $B_\alpha X = 0$ which is equivalent to $PA_\alpha = A_\alpha P$, for any $\alpha \in \{1, \dots, r\}$.

For $r = 1$ we have $g(Y, \xi)BX = 0$ (where $B = PA - AP$) and for $Y = \xi$ we have $g(\xi, \xi)BX = 0$. But $g(\xi, \xi) = q + pa - a^2 \neq 0$ (M is non-invariant submanifold) and we obtain $BX = 0$, for any $X \in \mathcal{X}(M)$, which is equivalent to $PA = AP$. \square

Remark 5.5. If M is a non-invariant totally umbilical (or totally geodesic) n -dimensional submanifold of codimension $r \geq 1$ in a locally metallic Riemannian manifold $(\overline{M}, \overline{g}, J)$ such that the normal connection ∇^\perp vanishes identically, then the Σ -metallic Riemannian structure induced on the submanifold M is normal.

Acknowledgements. The authors want to express their gratitude to the referees of this paper.

References

- [1] M. Crasmareanu and C.E. Hretcanu, *Golden differential geometry*, Chaos Solitons Fractals **38** no. 5 (2008), 1229–1238.
- [2] M. Crasmareanu, C.E. Hretcanu and M.I. Munteanu, *Golden- and product-shaped hypersurfaces in real space forms*, Int. J. Geom. Methods Mod. Phys. 10 no. 4 (2013), 1320006, 9 pp.
- [3] S.I. Goldberg and K. Yano, *Polynomial structures on manifolds*, Kodai Math. Sem. Rep. **22** (1970), 199–218.
- [4] S.I. Goldberg and N.C. Petridis, *Differentiable solutions of algebraic equations on manifolds*, Kodai Math. Sem. Rep. **25** (1973), 111–128.
- [5] C.E. Hretcanu, *Induced structures on spheres and product of spheres*, An. Stiint. Univ. Al. I. Cuza Iasi Mat. (N.S.), **54**, 1 (2008), 39–50.
- [6] C.E. Hretcanu and M. Crasmareanu, *On some invariant submanifolds in a Riemannian manifold with Golden structure*, An. Stiint. Univ. Al. I. Cuza Iasi. Mat. (N.S.) **53** suppl. 1 (2007), 199–211.
- [7] C.E. Hretcanu and M.C. Crasmareanu, *Applications of the Golden ratio on Riemannian manifolds*, Turkish J. Math. **33**, 2, (2009), 179–191.
- [8] C.E. Hretcanu and M. Crasmareanu, *Metallic structures on Riemannian manifolds*, Rev. Un. Mat. Argentina **54**, 2, (2013), 15–27.
- [9] G. Pitis, *On some submanifolds of a locally product manifold*, Kodai Math. J. **9**, 3, (1986), 327–333.
- [10] V.W. de Spinadel, *The metallic means family and forbidden symmetries*, Int. Math. J. **2**, 3, (2002), 279–288.
- [11] K. Yano and M. Kon, *Structures on Manifolds*, World Scientific, Singapore, Series in Pure Mathematics 3, 1984.

Authors' addresses:

Cristina Elena Hretcanu
Ștefan cel Mare University of Suceava, Romania.
E-mail: criselenab@yahoo.com

Adara Monica Blaga
West University of Timisoara, Romania.
E-mail: adarablaga@yahoo.com